

Your naloxone purchase should include:

- At least 2 doses 0.4 mg/ml naloxone
- At least 2 retractable VanishPoint® syringes

What is Naloxone and Why Do We Need It?

Naloxone is an antidote to opioid overdose. Taking too much of opioid drugs (like morphine, oxycodone, methadone, heroin, or fentanyl) can slow down or stop breathing. Naloxone restores normal breathing and consciousness within 1 to 5 minutes of injection. Giving naloxone can prevent death or brain damage from lack of oxygen during an opioid overdose. Naloxone can be injected through clothing into the big muscle of the thigh, arm or buttocks.

Signs of Opioid Overdose



Not moving and can't be woken



Slow or not breathing



Blue lips and nails



Choking, gurgling sounds or snoring



Cold or clammy skin



Tiny pupils

Overdose is a medical emergency - CALL 9-1-1 | Tell attendant: Person is not responsive and not breathing.

HealthLink BC: 8-1-1 **BC Drug & Poison Information Centre:** 1-800-567-8911

SAVE ME Steps

Follow the **SAVE** ME steps below to respond



If the person must be left unattended at any time, put them in the recovery position



Unresponsive?

Call 911



Check and open





VENTILATE

1 breath every

5 seconds



EVALUATE

Breathing?



MEDICATION

1 dose of

naloxone



EVALUATE & SUPPORT

Wait 5 minutes. Another dose?

How to Give Breaths



Tilt head back to open the airway

Give 1 breath

every 5 seconds



Pinch nose. Cover mouth with mask or clothing

Breathing can slow down during an overdose but the heart is probably still beating - Give breaths to keep them alive

How to Use Naloxone










Tap ampoule to send all liquid to the bottom

Push top away from you to snap open the ampoule



Pull plunger to draw up liquid





Press plunger all the way down to trigger safety (retraction)

Developed with the BC Centre for Disease Control

College of Pharmacists of British Columbia

bcpharmacists.org/naloxone